# SIGN AND RETURN

# Student-athlete & Parent/Legan Guardian Concussion Statement

Must be signed and returned to school or cummunity youth	athletic activity prior to
participation in practice or play	

Student-Athlete Name: \_\_\_\_\_ Grade: \_\_\_\_\_ Parent/Legal Guardian Name(s): After reading the information sheet, I am aware of the following information: Student-Parent/Legal Athlete Guardian Initials Initials A concussion is a brain injury which should be reported to my parents, my coach(es) or a medical professional if one is available A concussion cannot be "seen." Some symptons might be present right away. Other symptoms can show up hours or days after an injury I will tell my parents, my coach and/or a medical professional about my injuries and illnesses. N/A I will not return to play in a game or practice if a hit to my head or body causes any concussion-related symptoms. N/A I will/mychild will need written permission from a health care provider\* to return to play or practice after a consussion Most concussions take days or weeks to get better. A more serious concussion can last for months or longer. After a bump, blow, or jolt to the head or body an athlete should receive immediate medical attention if there are any danger signs such as loss of consciousness, repeated vomiting or a headache that gets worse. After a concussion, the brain needs time to heal. I understand that I am/my child is much more likely to have another concussion or more serious brain injury if return to play or practice occurs before the concussion symptoms go away sometimes repeat concussion can cause serious and long-lasting problems and even death. I have read the concussion symptoms on the Concussion Information Sheet \*Health care provider means a Tennessee licensed medical doctor, osteropathic physician or a clinical Neuropsychologist with consussion traing **Signature of Student-Athlete** Date

Date

Signature of Parent/Legal guardian

# (C(0)N(CILISISII(0)N)

### INFORMATION AND SIGNATURE FORM

# FOR STUDENT-ATHLETES & PARENTS/LEGAL GUARDIANS

(Adapted from CDC "Heads Up Concussion in Youth Sports")

Public Chapter 148, effective January 1, 2014, requires that school and community organizations sponsoring youth athletic activities establish guidelines to inform and educate coaches, youth athletes and other adults involved in youth athletics about the nature, risk and symptoms of concussion/head injury.

Read and keep this page.

Sign and return the signature page.

A concussion is a type of traumatic brain injury that changes the way the brain normally works. A concussion is caused by a bump, blow or jolt to the head or body that causes the head and brain to move rapidly back and forth. Even a "ding," "getting your bell rung" or what seems to be a mild bump or blow to the head can be serious.

### Did You Know?

- Most concussions occur without loss of consciousness.
- Athletes who have, at any point in their lives, had a concussion have an increased risk for another concussion.
- Young children and teens are more likely to get a concussion and take longer to recover than adults.

### WHAT ARE THE SIGNS AND SYMPTOMS OF CONCUSSION?

Signs and symptoms of concussion can show up right after the injury or may not appear or be noticed until days or weeks after the injury.

If an athlete reports one or more symptoms of concussion listed below after a bump, blow or jolt to the head or body, s/he should be kept out of play the day of the injury and until a health care provider\* says s/he is symptom-free and it's OK to return to play.

SIGNS OBSERVED BY COACHING STAFF	SYMPTOMS REPORTED BY ATHLETES
Appears dazed or stunned	Headache or "pressure" in head
	Render of the control
Forgets an instruction	Balance problems or dizziness
ોર્ડ પ્રાપ્તાનહીં શ્રી વ્યવસાર જુના કરાઈ છે એમાં આવેલા 🚡	Darble or bling restore &
Moves clumsily	Sensitivity to light
Annayyəta innərmin kalınıylıy 🍙 . 🥻 🥻	
Loses consciousness, even briefly	Feeling sluggish, hazy, foggy or groggy
Shows interest basic value of the same property of the contraction of	Congenitation originally problems
Can't recall events prior to hit or fall	Confusion
(បំរាស ស្ត្រីស្ត្រីអាស្ត្រស្វីអាស្ត្រីស្ត្រីស្រុស អាស៊ីលា ផ្លេង!	j dugingi Kedingaghi o Kedingawa . 😿 🧱

<sup>\*</sup>Health care provider means a Tennessee licensed medical doctor, osteopathic physician or a clinical neuropsychologist with concussion training

### **CONCUSSION DANGER SIGNS**

In rare cases, a dangerous blood clot may form on the brain in a person with a concussion and crowd the brain against the skull. An athlete should receive immediate medical attention after a bump, blow or jolt to the head or body if s/he exhibits any of the following danger signs:

- One pupil larger than the other
- Is drowsy or cannot be awakened
- A headache that not only does not diminish, but gets worse
- Weakness, numbness or decreased coordination
- Repeated vomiting or nausea
- Slurred speech
- Convulsions or seizures
- Cannot recognize people or places
- Becomes increasingly confused, restless or agitated
- Has unusual behavior
- Loses consciousness (even a brief loss of consciousness should be taken seriously)

# WHY SHOULD AN ATHLETE REPORT HIS OR HER SYMPTOMS?

If an athlete has a concussion, his/her brain needs time to heal. While an athlete's brain is still healing, s/he is much more likely to have another concussion. Repeat concussions can increase the time it takes to recover. In rare cases, repeat concussions in young athletes can result in brain swelling or permanent damage to their brains. They can even be fatal.

### Remember

Concussions affect people differently. While most athletes with a concussion recover quickly and fully, some will have symptoms that last for days, or even weeks. A more serious concussion can last for months or longer.

# WHAT SHOULD YOU DO IF YOU THINK YOUR ATHLETE HAS A CONCUSSION?

If you suspect that an athlete has a concussion, remove the athlete from play and seek medical attention. Do not try to judge the severity of the injury yourself. Keep the athlete out of play the day of the injury and until a health care provider\* says s/he is symptom-free and it's OK to return to play.

Rest is key to helping an athlete recover from a concussion. Exercising or activities that involve a lot of concentration such as studying, working on the computer or playing video games may cause concussion symptoms to reappear or get worse. After a concussion, returning to sports and school is a gradual process that should be carefully managed and monitored by a health care professional.

\* Health care provider means a Tennessee licensed medical doctor, osteopathic physician or a clinical neuropsychologist with concussion training.

# Athlete/Parent/Guardian Sudden Cardiac Arrest Symptoms and Warning Signs Information sheet and Acknowledgement of Receipt and Review Form

#### What is sudden cardiac arrest?

Sudden cardiac arrest (SCA) is when the heart stops beating, suddenly and unexpectedly. When this happens, blood stops flowing to the brain and other vital organs. SCA doesn't just happen to adults; it takes the lives of students, too. However, the causes of sudden cardiac arrest in students and adults can be different. A youth athlete's SCA will likely result from an inherited condition, while an adult's SCA may be caused by either inherited or lifestyle issues. SCA is NOT a heart attack. A heart attack may cause SCA, but they are not the same. A heart attack is caused by a blockage that stops the flow of blood to the heart. SCA is a malfunction in the heart's electrical system, causing the heart to suddenly stop beating.

### How common is sudden cardia arrest in the United States?

SCA is the #1 cause of death for adults in this country. There are about 300,000 cardiac arrests outside hospitals each year. About 2,000 patients under 25 die of SCA each year. It is the #1 cause of death for student athletes.

#### Are there warning signs?

Although SCA happens unexpectedly, some people may have signs or symptoms, such as:

- Fainting or seizures during exercise;
- Unexplained shortness of breath;
- Dizziness;
- Extreme fatigue;
- Chest pains; or
- Racing heart.

These symptoms can be unclear in athletes, since people often confuse these warning signs with physical exhaustion. SCA can be prevented if the underlying causes can be diagnosed and treated.

### What are the risks of practicing or playing after experiencing these symptoms?

There are risks associated with continuing to practice or play after experiencing these symptoms. When the heart stops, so does the blood that flows to the brain and other vital organs. Death or permanent brain damage can occur in just a few minutes. Most people who experience SCA die from it.

Public Chapter 325 – the Sudden Cardiac Arrest Prevention Act

The act is intended to keep youth athletes safe while practicing or playing. The requirements of the act are:

All youth athletes and their parents or guardians must read and sign this form. It must be returned to the school before participation in any athletic activity. A new form must be signed and returned each school year.

	Signature of Parent/Guardian			DATE
Signature of Student-Athlete		ent-Athlete	Print Student-Athlete's Name	DATE
- I have re	the process to the pr	ractice or competition during wastent with sudden cardiac arrested returning to practice or play is essee licensed medical doctor cated return to practice or play	in an athletic activity, the athlete must be or an osteopathic physician. Clearance to	toms evaluated by a
	(ii) (iii) (iv) (v)	Chest pains; Dizziness Racing heart rate; or Extreme fatigue; and		
		• •	h athlete who passes out or faints while p any of the following symptoms reath:	articipating in

V =